

CACI Contra-Indications

Never Treat

Pacemaker - Microcurrent could disturb heart rate.

Epilepsy - Microcurrent may cause a seizure.

Tumors - Increased blood flow can stimulate growth.

Pregnancy - If halfway through course, stop immediately.

Thrombosis / Phlebitis - Microcurrent could disturb / cause clotting.

GP Letter

Heart Conditions - i.e Angina / Coronary Heart Disease

Diabetes - Okay if diet is controlled, if insulin dependent retrieve a GP note.

Severe Muscular Conditions - i.e Muscular Dystrophy

All Forms of Cancer - either 5 years in remission, or if in remission provide a consultant / GP letter.

Avoid Area

Botox & Thread Lift - Avoid the area for 3 months, Botox may disperse if current is applied over it.

Injectables - Avoid area for 2 weeks due to discomfort.

Inflammation / Skin Diseases - Avoid area due to cross contaminations.

Metal Pins / Plates

The Contraceptive Coil

Under & Over Active Thyroid - Avoid area

IPL / Laser - Avoid for 3-5 days

Other

High Blood Pressure Med & Anti-Depressants - Results may vary due to muscle relaxants in meds.

Multiple Sclerosis - Results may be limited due to nerve damage caused by MS.

Retin A - Wait 3-6 months after completion.

Roaccutane / Accutane - Wait 6 months after completion.

Dental Implants - May experience stronger metallic taste in mouth.

Allergy to rubber / metal / silicone - CACI Probes are stainless steel, body and facial pads are rubber based and coated in silicone.